




2023 State of the Ecosystem

















Mid-Atlantic



NOAA
FISHERIES

Performance Relative to Fishery Management Objectives

Trends and status of indicators related to broad ecosystem-level fishery management objectives, with implications for the Mid-Atlantic Fishery Management Council (MAFMC)

OBJECTIVE (Indicator)	TREND	CURRENT STATUS	IMPLICATIONS
Seafood production (total and MAFMC managed landings)	 Decline	 Below long term average	Commercial landings are at the lowest point observed, driven by recent declines in species not managed by the Mid-Atlantic Council. Recreational harvest is declining due to multiple drivers. COVID-19 likely exacerbated existing trends, but impacts are not uniform across fisheries. Biomass trends within the ecosystem continue to be stable. Climate indicators continue to exceed historical bounds, which affects stock distributions and will generate other ecosystem changes.
Commercial profits	 Decline	 Below long term average	Regional commercial revenue is the lowest that has been observed, driven in part by managed clam species. Falling prices are almost universal and due to market dynamics including COVID-19 impacts. Monitor climate risks to surfclams and ocean quahogs.
Recreational opportunities (effort and fleet diversity)	 No trend	 Near long term average	Recreational effort shows no long term trend and is near average, but fleet diversity is decreasing because of a shift away from party/charter to shore-based fishing. This shift results in a decreased range of recreational fishing opportunities. Shore-based anglers will have access to different species/sizes of fish than vessel-based anglers.
	 Decline	 Below long term average	
Stability (fishery and ecosystem diversity maintained over time)	 No trend	 Near long term average	Commercial: Fleet diversity metrics suggest stable capacity to respond to the current range of fishing opportunities. Recreational: Species catch diversity has been maintained by a different set of species over time and continues to be above the long-term mean. Ecosystem: Adult fish diversity indices are stable, but several climate and oceanography metrics are changing and should be monitored as warning signs for potential regime shift or ecosystem restructuring.
	 Mixed trends	 Near long term average	
Social and cultural (community fishery engagement, reliance, and environmental justice vulnerability)	Status only indicator	Environmental justice status for top commercial and recreational communities	These indicators are used to identify top fishing communities and those with environmental justice concerns based on 2020 data. Highlighted communities may be vulnerable to changes in fishing patterns due to regulations and/or climate change. When any of these communities also experience environmental justice issues, they may have lower ability to successfully respond/adapt to change. The top Mid Atlantic recreational communities changed between 2019 and 2020.
Protected species (coastwide bycatch, population numbers, mortalities)	 Mixed trends	 Meeting objectives	Mixed bycatch trends through 2021 are related to fishery management, shifts in population distribution combined with fishery shifts, and population increase for seals. Recent bycatch data is uncertain. Population drivers for North Atlantic Right Whales (NARW) include combined fishery interactions/vessel strikes, distribution shifts, and copepod availability. Unusual mortality events continue for 3 large whale species.
	 Decline	 Below long term average	

Risks to Meeting Fishery Management Objectives

Climate and Ecosystem Productivity Risks

Climate change, most notably ocean warming and changes in the Gulf Stream, continue to affect the Mid-Atlantic ecosystem:

- 2022 was among the warmest years on record in the North Atlantic, with both long term surface and bottom warming observed in the Mid-Atlantic.
- The Gulf Stream is becoming less stable and moving further north, which can affect the physics, chemistry, and biology of the Northeast Shelf.
- The cold pool is becoming warmer, smaller, and shorter in duration, which affects habitat for multiple federally managed species.
- Ocean acidification in western Long Island Sound, nearshore to mid-shelf waters of the Mid-Atlantic Bight off the coast of New Jersey, and in waters > 1000 meters may impact organisms.
- Above average early winter and late fall phytoplankton blooms were observed in the Mid-Atlantic, but larger phytoplankton concentrations were below average in early fall.
- The value of Chesapeake Bay habitat for fishes is changing. Several finfish species, including summer flounder, show relative decline in Chesapeake Bay habitat usage. There is evidence that suitable habitat for juvenile summer flounder has declined between 47% and 64% since 1996.
- Shifts in species distribution are being observed across many managed fish and marine mammal species, complicating regional management by changing fishing patterns and risks.
- Fish condition was mixed in 2022, and fish productivity is declining for many managed species.



Other Ocean Uses: Offshore Wind Risks



More than 31 offshore wind development projects are proposed for construction on the Northeast shelf, covering more than 2.4 million acres by 2030. Additional large areas are being considered. In existing and proposed leases of the Northeast:

- 1–34% of port revenue from fisheries currently comes from areas proposed for offshore wind development. Some of these port communities score medium-high to high in environmental justice concerns and gentrification vulnerability.
- Up to 17% of annual commercial landings and revenue for Mid-Atlantic managed species occur in lease areas and may shift to other areas.
- Development at different scales will affect species differently, negatively affecting species that prefer soft bottom habitat while potentially benefiting species that prefer hard structured habitat.
- Planned wind areas overlap with important right whale foraging habitats, and altered local oceanography could affect right whale prey availability. Development also brings increased vessel strike risk and the potential impacts of pile driving noise.
- Scientific surveys are key to understanding the impacts of climate change and other drivers on managed species, and inform management advice. Planning for impacts to scientific surveys is in progress.
- Current plans for rapid buildout in a patchwork of areas would spread the impacts differentially throughout the region.

Characterizing Ecosystem Change

Multiple System Drivers

The Northeast shelf ecosystem is changing, which is affecting the services that the ecosystem provides. To illustrate how multiple factors are driving change in this complex ecosystem, we are using three overarching concepts: multiple system drivers, regime shifts, and ecosystem reorganization. Societal, biological, physical, and chemical factors are the **multiple system drivers** that influence marine ecosystems through a variety of different pathways.



Regime Shift

These drivers affect fishery management objectives such as seafood production and recreational opportunities, as well as other ecosystem services we derive from the ocean. Changes in the multiple drivers can lead to **regime shifts**—large, abrupt and persistent changes in the structure and function of an ecosystem. Regime shifts and changes in how multiple system drivers interact can result in **ecosystem reorganization** as species and humans respond and adapt to the new environment.

Introduction

About This Report

This report is for the Mid-Atlantic Fishery Management Council (MAFMC). The purpose of this report is to synthesize ecosystem information to allow the MAFMC to better meet fishery management objectives, and to update the MAFMC’s Ecosystem Approach to Fishery Management (EAFM) risk assessment. The major messages of the report are synthesized on pages 1 and 2, and synthesis themes are illustrated on page 3. The information in this report is organized into two sections; **performance measured against ecosystem-level management objectives** (Table 1), and potential **risks to meeting fishery management objectives** (climate change and other ocean uses).

Report structure

The two main sections contain subsections for each management objective or potential risk. Within each subsection, we first review indicator trends, and the status of the most recent data year relative to a threshold (if available) or relative to the long-term average. Second, we synthesize results of other indicators and information to outline potential implications for management (i.e., connecting indicator(s) status to management and why an indicator(s) is important). For example, if there are multiple drivers related to an indicator trend, which drivers may be more or less supported by current information, and which, if any, can be affected by management action(s)? Similarly, which risk indicators warrant continued monitoring to evaluate whether regime shifts or ecosystem reorganization are likely? We emphasize that these implications are intended to represent testable hypotheses at present, rather than “answers,” because the science behind these indicators and syntheses continues to develop.

A glossary of terms¹, detailed technical methods documentation², and indicator data³ are available online. The details of standard figure formatting (Fig. 57a), categorization of fish and invertebrate species into feeding guilds (Table 3), and definitions of ecological production units (EPUs, including the Mid-Atlantic Bight, MAB; Fig. 57b) are provided at the end of the document.

Table 1: Ecosystem-scale fishery management objectives in the Mid-Atlantic Bight

Objective categories	Indicators reported
Provisioning and Cultural Services	
Seafood Production	Landings; commercial total and by feeding guild; recreational harvest
Profits	Revenue decomposed to price and volume
Recreation	Angler trips; recreational fleet diversity
Stability	Diversity indices (fishery and ecosystem)
Social & Cultural	Community engagement/reliance and environmental justice status
Protected Species	Bycatch; population (adult and juvenile) numbers, mortalities
Supporting and Regulating Services	
Biomass	Biomass or abundance by feeding guild from surveys
Productivity	Condition and recruitment of managed species, primary productivity
Trophic structure	Relative biomass of feeding guilds, zooplankton
Habitat	Estuarine and offshore habitat conditions

Performance Relative to Fishery Management Objectives

In this section, we examine indicators related to broad, ecosystem-level fishery management objectives. We also provide hypotheses on the implications of these trends—*why* we are seeing them, what’s driving them, and potential or observed regime shifts or changes in ecosystem structure. Identifying multiple drivers, regime shifts, and potential changes to ecosystem structure, as well as identifying the most vulnerable resources, can help managers determine whether we can do anything differently to meet objectives and how to prioritize for upcoming issues/risks.

¹<https://noaa-edab.github.io/tech-doc/glossary.html>

²<https://NOAA-EDAB.github.io/tech-doc>

³<https://github.com/NOAA-EDAB/ecodata>

Seafood Production

Indicators: Landings; commercial and recreational

This year, we present updated indicators for total commercial landings (all species, all uses, fleets from all nations), US seafood landings (species for human consumption landed by US fleets), and Council-managed US seafood landings (Mid-Atlantic Fishery Management Council (MAFMC) and jointly managed species landed by US fleets for human consumption). Total commercial landings (black) within the Mid-Atlantic have declined over the long term, and total US seafood landings are near their all time low. Because there is no long term trend in MAFMC managed US seafood landings, the decline in US seafood landings in the Mid-Atlantic region is likely driven by recent declines in species not managed by the Mid-Atlantic Council (Fig. 1).




Figure 1: Total commercial landings (black), total U.S. seafood landings (blue), and Mid-Atlantic managed U.S. seafood landings (red)

Landings by guild include all species and all uses, and are reported as total for the guild and the MAFMC managed species within the guild. As reported in previous years, landings of benthos presented a significant downward trend, primarily driven by surf clam and ocean quahog. However, total landings of planktivores is now also presenting a significant downward trend, primarily due to decreases in species not managed by the Mid-Atlantic Council (Atlantic herring and Atlantic menhaden; Fig. 2).




Figure 2: Total commercial landings (black) and MAFMC managed U.S. seafood landings (red) by feeding guild.

Total recreational harvest (retained fish presumed to be eaten) is down in the MAB (Fig. 3). Although harvest has increased from a historic low in 2018, it is still below the long term average.




Figure 3: Total recreational seafood harvest (millions of pounds) in the Mid-Atlantic region.

Recreational shark landings show an increase in pelagic sharks over the past decade, with a sharp decrease in 2018 - 2019 persisting through 2022 (Fig 4). This is likely influenced by regulatory changes implemented in 2018 intended to rebuild shortfin mako stocks. In 2021 the International Commission for the Conservation of Atlantic Tunas (ICCAT) finalized recommendations for a two-year retention ban for shortfin mako (ICCAT Rec.21-09), which will also affect total overall landings of pelagic sharks in coming years.




Figure 4: Recreational shark landings from Marine Recreational Information Program.

Aquaculture production is not yet included in total seafood landings, but we are working toward including it in future reports. Available aquaculture production of oysters for a subset of Mid-Atlantic states indicates a decline in recent years.⁴

Implications

Declining commercial (total and seafood) and recreational landings can be driven by many interacting factors, including combinations of ecosystem and stock production, management actions, market conditions (including COVID-19 disruptions), and environmental change. While we cannot evaluate all possible drivers at present, here we evaluate the extent to which stock status and system biomass trends may play a role.

Stock Status and Catch Limits Single species management objectives (1. maintaining biomass above minimum thresholds and 2. maintaining fishing mortality below overfishing limits) are being met for all but one MAFMC managed species, though the status of six stocks is unknown (Fig. 5). In addition, the status of Spiny dogfish and bluefish are based on 2022 research track assessments and are thus waiting for a management track update to finalize stock status.

⁴https://noaa-edab.github.io/ecodata/human_dimensions_MAB#Commercial; “Oyster Aquaculture” tab




Figure 5: Summary of single species status for MAFMC and jointly federally managed stocks (Spiny dogfish and both Goosefish). The dotted vertical line is the target biomass reference point of B_{msy} . The dashed lines are the management thresholds of one half B_{msy} (vertical) or F_{msy} (horizontal). Stocks in red are below the biomass threshold (overfished) and have fishing mortality above the limit (subject to overfishing), stocks in green are above the biomass threshold but have fishing mortality above the limit. Remaining stocks have fishing mortality within limits: stocks in orange are above the biomass threshold but below the biomass target, and stocks in purple are above the biomass target.

Stock status affects catch limits established by the Council, which in turn may affect landings trends. Summed across all MAFMC managed species, total Acceptable Biological Catch or Annual Catch Limits (ABC or ACL) have been relatively stable 2012-2020 (Fig. 6). Although these figures have not been updated with 2021 data, we do not expect a single year's update to change the narrative. The recent total ABC or ACL is lower relative to 2012-2013, with much of that decrease due to declining Atlantic mackerel ABC. This is true even with the addition of blueline tilefish management in 2017 contributing an additional ABC and ACL to the total 2017-2020, due to that fishery's small relative size.




Figure 6: Sum of catch limits across all MAFMC managed commercial (C) and recreational (R) fisheries.

Nevertheless, the percentage caught for each stock’s ABC/ACL suggests that these catch limits are not generally constraining as most species are well below the 1/1 ratio (Fig. 7). Therefore, stock status and associated management constraints are unlikely to be driving decreased landings for the majority of species.




Figure 7: Catch divided by ABC/ACL for MAFMC managed fisheries. High points = Recreational Black Sea Bass. Red line indicates the median ratio across all fisheries.

System Biomass Although aggregate biomass trends derived from scientific resource surveys are mostly stable in the MAB, spring piscivores, spring benthivores, and fall benthos show long-term increases (Fig. 8). While managed species make up varying proportions of aggregate biomass, trends in landings are not mirroring shifts in the overall trophic structure of survey-sampled fish and invertebrates. Therefore, major shifts in feeding guilds or ecosystem trophic structure are unlikely to be driving the decline in landings.




Figure 8: Spring (left) and fall (right) surveyed biomass in the Mid-Atlantic Bight. Data from the NEFSC Bottom Trawl Survey are shown in black, with the nearshore NEAMAP survey shown in red. The shaded area around each annual mean represents 2 standard deviations from the mean.

Effect on Seafood Production Stock status is above the minimum threshold for all but one stock, and aggregate biomass trends appear stable, so the decline in commercial seafood landings is most likely driven by market dynamics affecting the landings of surfclams and ocean quahogs, as landings have been below quotas for these species. The long term decline in total planktivore landings is largely driven by Atlantic menhaden fishery dynamics, including a consolidation of processors leading to reduced fishing capacity between the 1990s and mid-2000s.

Climate change also seems to be shifting the distribution of surfclams and ocean quahogs, resulting in areas with overlapping distributions and increased mixed landings. Given the regulations governing mixed landings, this could become problematic in the future and is currently being evaluated by the Council.

The decline in recreational seafood harvest stems from other drivers. Some of the decline, such as that for recreational shark landings, is driven by management intended to reduce fishing mortality on mako sharks. However, NOAA Fisheries' Marine Recreational Information Program survey methodology was updated in 2018, so it is unclear whether the record-low landings for species other than sharks in 2018 are driven by changes in fishing behavior or the change in the survey methodology. Nevertheless, the recreational harvest seems to be stabilizing at a lower level than historical estimates.

Other environmental changes require monitoring as they may become important drivers of landings in the future:

- Climate is trending into uncharted territory. Globally, 2022 was among the warmest years on record⁵ (see [Climate Risks](#) section).
- Stocks are shifting distribution, moving towards the northeast and into deeper waters throughout the Northeast US Large Marine Ecosystem (Fig. 9).




Figure 9: Aggregate species distribution metrics for species in the Northeast Large Marine Ecosystem.

- Some ecosystem composition and production changes have been observed (see [Stability](#) section).
- Some fishing communities are affected by environmental justice vulnerabilities (see [Environmental Justice and Social Vulnerability](#) section).

Commercial Profits

Indicators: revenue (a proxy for profits)

Total commercial revenues (black) within the Mid-Atlantic and Mid-Atlantic managed species revenue both present long-term declining trends. Total revenue is at, and revenue from Mid-Atlantic managed species is near, an all-time low (Fig. 10).

⁵<https://www.climate.gov/news-features/understanding-climate/climate-change-global-temperature>




Figure 10: Revenue for the for the Mid-Atlantic region: total (black) and from MAFMC managed species (red).

Revenue earned by harvesting resources is a function of both the quantity landed of each species and the prices paid for landings. Beyond monitoring yearly changes in revenue, it is even more valuable to determine what drives these changes: harvest levels, the mix of species landed, price changes, or a combination of these. The Bennet Indicator decomposes revenue change into two parts, one driven by changing quantities (volumes), and a second driven by changing prices.

Total revenue trends, decomposed to price and volume indicators (Fig. 11), mirror price and volume indicator trends for the benthos (clams; orange in Fig. 12) group, especially over the past decade. However, of note is that only piscivore volume is up across species guilds for either prices or volume when compared to the 2015 benchmark year.




Figure 11: Revenue change from the 2015 values in dollars (black), Price (PI), and Volume Indicators (VI) for commercial landings in the Mid-Atlantic Bight.




Figure 12: Total component value in dollars (black) for commercial landings in the Mid-Atlantic Bight.

Implications

In a similar manner to seafood landings, the results here are driven in large part by market dynamics affecting the landings of surfclams and ocean quahogs, as landings have been below quotas for these species. Changes in other indicators, particularly those driving landings and those related to climate change, require monitoring as they may become important drivers of revenue in the future; for example:

- Surfclams and ocean quahogs are sensitive to warming ocean temperatures and ocean acidification.
- Acidification levels in surfclam summer habitat are approaching, but not yet at, levels affecting surfclam growth (see [Climate Risks](#) section).

Recreational Opportunities

Indicators: Angler trips, fleet diversity

Recreational effort (angler trips) in 2021 is around the long-term average (Fig. 13). However, recreational fleet diversity (i.e., effort by shoreside, private boat, and for-hire anglers) has declined over the long term (Fig. 14).




Figure 13: Recreational effort in the Mid-Atlantic.




Figure 14: Recreational fleet effort diversity in the Mid-Atlantic.

Implications

While the overall number of recreational opportunities in the MAB is above the long-term average, the continuing decline in recreational fleet effort diversity suggests a potentially reduced range of recreational fishing options, despite the slight increase in this indicator’s value between 2020 and 2021.

The downward effort diversity trend is driven by party/charter contraction (2% currently), and a shift toward shorebased angling, which currently makes up 61% of angler trips.. Effort in private boats remains stable at around 37% of trips.

Changes in recreational fleet diversity can be considered when managers seek options to maintain recreational opportunities. Shore anglers will have access to different species than vessel-based anglers, and when the same species is accessible both from shore and from a vessel, shore anglers typically have access to smaller individuals. Many states have developed shore-based regulations where the minimum size is lower than in other areas and sectors to maintain opportunities in the shore angling sector.

Stability

Indicators: fishery fleet and catch diversity, ecological component diversity

While there are many potential metrics of stability, we use diversity indices as a first check to evaluate overall stability in fisheries and ecosystems. In general, diversity that remains constant over time suggests a similar capacity to respond to change over time. A significant change in diversity over time does not necessarily indicate a problem or an improvement, but does indicate a need for further investigation. We examine commercial fleet and species catch diversity, and recreational species catch diversity (with fleet effort diversity discussed above), and diversity in zooplankton, and larval and adult fishes.

Fishery Diversity Diversity estimates have been developed for fleets landing managed species, and species landed by commercial vessels with Mid-Atlantic permits. A fleet is defined here as the combination of gear type (Scallop Dredge, Other Dredge, Gillnet, Hand Gear, Longline, Bottom Trawl, Midwater Trawl, Pot, Purse Seine, or Clam Dredge) and vessel length category (less than 30 ft, 30 to 50 ft, 50 to 75 ft, 75 ft and above). Commercial fishery fleet count and fleet diversity have been stable over time in the MAB, with current values near the long-term average (Fig. 15). This indicates similar commercial fleet composition and species targeting opportunities over time.




Figure 15: Commercial fleet count and diversity in the Mid-Atlantic.

Commercial fisheries are relying on fewer species relative to the mid-90s, and current species revenue diversity is near the historical low point (Fig. 16). Although with precedent, the drop between 2020 and 2021 is relatively large.




Figure 16: Species revenue diversity in the Mid-Atlantic.

As noted above, recreational fleet effort diversity is declining (Fig. 14), so this metric suggests an unstable range of recreational fishing opportunities. However, recreational species catch diversity has no long term trend so is considered stable, and has been at or above the long term average in 7 of the last 10 years (Fig. 17).




Figure 17: Diversity of recreational catch in the Mid-Atlantic.

Ecological Diversity Ecological diversity indices show mixed trends. Zooplankton diversity is increasing in the MAB (Fig. 18). Larval fish diversity shows no trend, and high interannual variability with 2021 values at the mean. Adult fish diversity is measured as the expected number of species in a standard number of individuals sampled from the NEFSC bottom trawl survey. There is no vessel correction for this metric, so indices collected aboard the research vessel Albatross IV (up to 2008) and research vessel Bigelow (2009-2021) are calculated separately. Despite this, adult fish diversity indices appear stable over time, with current values within one standard deviation from most historic estimates (Fig. 19).




Figure 18: Zooplankton diversity in the Mid-Atlantic Bight, based on Shannon diversity index.




Figure 19: Adult fish diversity in the Mid-Atlantic Bight, based on expected number of species. Results from survey vessels Albatross and Bigelow are reported separately due to catchability differences.

Implications

Fleet diversity indices are used by the MAFMC to evaluate stability objectives as well as risks to fishery resilience and maintaining equity in access to fishery resources [1].

Stability in commercial fleet diversity metrics suggests stable capacity to respond to the current range of fishing opportunities. However, commercial species diversity is relatively low, indicating substantial changes in fishing activities even as the fleet composition sees relative stability.

Declining recreational fleet effort diversity, as noted [above](#), indicates that the party/charter boat sector continues to contract, with shoreside angling becoming more important, as a percentage of recreational angler trips.

Stability in recreational species catch diversity has been maintained by a different set of species over time. A recent increase in Atlantic States Marine Fisheries Commission (ASMFC) and South Atlantic Fishery Management Council (SAFMC) managed species in recreational catch is helping to maintain diversity in the same range that MAFMC and New England Fishery Management Council (NEFMC) species supported in the 1990s.

Ecological diversity indices can provide insight into ecosystem structure. Changes in ecological diversity over time may indicate altered ecosystem structure with implications for fishery productivity and management [2]. Stable adult fish diversity indicates the same overall number and evenness over time, but doesn't rule out species substitutions (e.g., warm-water replacing cold-water). In addition, the change in survey vessels complicates interpretation of long-term fish diversity trends.

In the MAB, existing diversity indicators suggest overall stability in the fisheries and ecosystem components examined. However, declining recreational fleet diversity suggests a potential loss in the range of recreational fishing opportunities. Increasing zooplankton diversity (due to increases in abundance of several taxa and stable or declining dominance of an important copepod species) suggests a shift in the zooplankton community that warrants continued monitoring to determine if managed species are affected. In addition, the species diversity in landings warrants continued attention given its relatively low value and large year over year decline.

Environmental Justice and Social Vulnerability

Indicators: Environmental Justice and Social Vulnerability in commercial and recreational fishing communities

Social vulnerability measures social factors that shape a community's ability to adapt to change. A subset of these factors can be used to assess potential environmental justice issues. Environmental Justice is defined in Executive Order 12898 as federal actions intended to address disproportionately high and adverse human health and environmental effects of federal actions on minority and low-income populations. Three of the existing NOAA Fisheries Community Social Vulnerability Indicators (CSVIs), the Poverty Index, Population Composition Index, and Personal Disruption Index, can be used for mandated Environmental Justice analysis⁶.

Commercial fishery engagement measures the number of permits and dealers, and pounds and value landed in a community, while reliance expresses these numbers based on the level of fishing activity relative to the total population of a community. Recreational fishery engagement measures shore, private vessel, and for-hire fishing effort while reliance expresses these numbers based on fishing effort relative to the population of a community.

In 2022, we reported the top ten most engaged, and top ten most reliant commercial and recreational fishing communities and their associated environmental justice vulnerability based on 2019 data. Here we apply the same selection standard for top ten fishing communities for both sectors using 2020 data, and again examine the environmental justice vulnerability in this updated set of communities. Changes in fishing activity between years changed community engagement and reliance rankings, and changes in vulnerability indicators changed environmental justice vulnerability scores.

Communities plotted in the upper right section of Fig.20 scored high for both commercial engagement and reliance using both 2019 and 2020 data, including Cape May and Barnegat Light, NJ, and Reedville, VA. Communities that ranked medium-high or above for one or more of the environmental justice indicators in 2020 are highlighted in bright orange, including Newport News, VA; Atlantic City, NJ; and Beaufort, Columbia and Hobucken, NC. Hampton

⁶<https://www.fisheries.noaa.gov/national/socioeconomics/social-indicators-coastal-communities>

Bays/Shinnecock, NY ranked medium-high based on 2019 data but decreased to medium for its environmental justice vulnerability based on 2020 data reported here.




Figure 20: Commercial engagement, reliance, and environmental justice vulnerability for the top commercially engaged and reliant fishing communities in the Mid-Atlantic. Communities ranked medium-high or above for one or more of the environmental justice indicators are highlighted in bright orange. *Community scored high (1.00 and above) for both commercial engagement and reliance indicators.

Fig. 21 shows the detailed scores of the three environmental justice indicators for the same communities plotted in Fig.20. Communities are plotted clockwise in a descending order of commercial engagement scores from high to low, with the most highly engaged community, Cape May, NJ, listed on the top. Among the communities ranked medium-high or above for environmental justice vulnerability, Atlantic City, NJ scored high for all of the three environmental justice indicators. Columbia, NC scored high for the personal disruption index and the poverty index. Hobucken, NC scored high for the personal disruption index. Newport News, VA scored medium-high for the population composition index⁷. Beaufort, NC scored medium-high for the poverty index.

⁷Due to missing data, the Poverty Index is missing for Hobucken and Rodanthe, NC




Figure 21: Environmental justice indicators (Poverty Index, population composition index, and personal disruption index) for top commercial fishing communities in Mid-Atlantic. *Community scored high (1.00 and above) for both commercial engagement and reliance indicators.

Considerably more communities scored high for both recreational engagement and reliance based on 2020 data relative to 2019. Joining Barnegat Light, NJ in the upper right section are Babylon, NY, Nags Head, NC, Hatteras, NC, Stevensville, MD, Atlantic Highlands, NJ, Morehead City, NC, Montauk, NY, Point Pleasant Beach, NJ, Ocean City, MD, Point Lookout, NY, Manteo, NC, and Vandemere, NC. Fig.22. Communities that ranked medium-high or above for one or more of the environmental justice indicators are highlighted in bright orange, including Ocean City and Bivale, MD; Hatteras, Manteo, Vandemere, and Hobuken, NC.




Figure 22: Recreational engagement and reliance, and environmental justice vulnerability, for the top recreationally engaged and reliant fishing communities in the Mid-Atlantic. Communities ranked medium-high or above for one or more of the environmental justice indicators are highlighted in bright orange. *Community scored high (1.00 and above) for both recreational engagement and reliance indicators.

Fig. 23 orders communities clockwise in a descending order of recreational engagement scores from high to low, with the most highly engaged community, Babylon, NY, listed on the top. Among the communities with environmental

justice concerns, Hatteras and Vandemere, NC scored medium-high for personal disruption and poverty index. Ocean City, MD and Hobucken, NC scored medium-high for personal disruption index. Manteo, NC scored high for poverty index. Bivale, MD scored medium-high for population composition index⁸.




Figure 23: Environmental justice indicators (Poverty Index, population composition index, and personal disruption index) for top recreational fishing communities in Mid-Atlantic. *Community scored high (1.00 and above) for both recreational engagement and reliance indicators.

Both commercial and recreational fishing are important activities in Montauk, NY, Barnegat Light and Point Pleasant Beach, NJ, Hatteras and Hobucken, NC, meaning these communities may be impacted simultaneously by commercial and recreational regulatory changes. Among these communities, Hobucken scored high for the personal disruption index⁹. Hatteras scored medium-high for the personal disruption index and Poverty Index. Montauk, NY, Barnegat Light, Cape May and Point Pleasant Beach, NJ scored lower than medium-high for all of the three environmental justice indicators, indicating that environmental justice may not be a major concern in these communities at the moment based on the indicators analyzed.

Implications

There was an increase in recreational fishing activities in many of the top recreational communities from 2019 to 2020. This increase may be due to multiple factors including the recreational boating boom across the country¹⁰ and increasing interest in for-hire/charter recreational fishing trips as an preferred outdoor recreation activities and ways to social distance in response to the COVID-19 pandemic [3].

These plots provide a snapshot of the presence of environmental justice issues in the most highly engaged and most highly reliant commercial and recreational fishing communities in the Mid-Atlantic. These communities may be vulnerable to changes in fishing patterns due to regulations and/or climate change. When any of these communities are also experiencing social vulnerability including environmental justice issues, they may have lower ability to successfully respond to change.

Protected Species

Protected species include marine mammals protected under the Marine Mammal Protection Act, endangered and threatened species protected under the Endangered Species Act, and migratory birds protected under the Migratory Bird Treaty Act. In the Northeast U.S., endangered/threatened species include Atlantic salmon, Atlantic and shortnose sturgeon, all sea turtle species, and five baleen whales. Fishery management objectives for protected species generally focus on reducing threats and on habitat conservation/restoration. Here we report on the status

⁸Due to missing data, the Poverty Index is missing for Hobucken, NC, Bivalve and Georgetown, MD

⁹Due to missing data, the Poverty Index is missing for Hobucken, NC

¹⁰National Marine Manufacturers Association. 2021. U.S. Boat Sales Reached 13-Year High in 2020, Recreational Boating Boom to Continue through 2021. Available at: <https://www.nmma.org/press/article/23527>

of these actions as well as indicating the potential for future interactions driven by observed and predicted ecosystem changes in the Northeast U.S. Protected species objectives include managing bycatch to remain below potential biological removal (PBR) thresholds, recovering endangered populations, and monitoring unusual mortality events (UMEs).

Indicators: bycatch, population (adult and juvenile) numbers, mortalities

Average indices for both harbor porpoise (Fig. 24) and gray seal bycatch (Fig. 25) are below current PBR thresholds, meeting management objectives. However, the 2019 bycatch estimate for gray seals was highest in the time series.




Figure 24: Harbor porpoise average bycatch estimate for Mid-Atlantic and New England gillnet fisheries (blue) and the potential biological removal (red).

The annual estimate for gray seal bycatch has declined since 2019, in part driven by declining gillnet landings. In addition, estimates since 2019 have greater uncertainty stemming from low observer coverage since 2019. The rolling mean confidence interval remains just below the removal threshold.




Figure 25: Gray Seal average bycatch estimate for gillnet fisheries (blue) and and the potential biological removal (red).

The North Atlantic right whale population was on a recovery trajectory until 2010, but has since declined (Fig. 26). Reduced survival rates of adult females and diverging abundance trends between sexes have also been observed. It is estimated that there are fewer than 70 adult females remaining in the population.




Figure 26: Estimated North Atlantic right whale abundance on the Northeast Shelf.

North Atlantic right whale calf counts have generally declined after 2009 to the point of having zero new calves observed in 2018 (Fig. 27). However, since 2019, we have seen more calf births each year, with 20 births in 2022.




Figure 27: Number of North Atlantic right whale calf births, 1990 - 2021.

This year, the Unusual Mortality Event (UME) for North Atlantic right whales continued. Since 2017, the total UME right whale mortalities includes 35 dead stranded whales, 14 in the US and 21 in Canada. When alive but seriously injured whales (22) and sublethal injuries or ill whales (37) are taken into account, 94 individual whales are included in the UME. Recent research suggests that many mortalities go unobserved and the true number of mortalities are about three times the count of the observed mortalities [4]. The primary cause of death is “human interaction” from entanglements or vessel strikes¹¹.

A UME continued from previous years for humpback whales (2016-present); suspected causes include human interactions. A UME for both gray and harbor seals on the Maine coast was declared in June 2022 due to a high number of mortalities thought to be caused by highly pathogenic avian influenza virus. A UME for minke whales that began in 2017 remains open, but is pending closure as of January 2023¹².

Implications

Bycatch management measures have been implemented to maintain bycatch below PBR thresholds. The downward trend in harbor porpoise bycatch could also be due to a decrease in harbor porpoise abundance in US waters, reducing their overlap with fisheries, and a decrease in gillnet effort. The increasing trend in gray seal bycatch may be related to an increase in the gray seal population (U.S. pup counts).

The number of gray seals in U.S. waters has risen dramatically in the last three decades. Based on a survey conducted in 2016, the size of the gray seal population in the U.S. during the breeding season was approximately

¹¹<https://www.fisheries.noaa.gov/national/marine-life-distress/2017-2023-north-atlantic-right-whale-unusual-mortality-event>

¹²<https://www.fisheries.noaa.gov/national/marine-life-distress/active-and-closed-unusual-mortality-events>

27,000 animals, while in Canada the population was estimated to be roughly 425,000. The population in Canada is increasing at roughly 4% per year, and contributing to rates of increase in the U.S., where the number of pupping sites has increased from one in 1988 to nine in 2019. Mean rates of increase in the number of pups born at various times since 1988 at four of the more data-rich pupping sites (Muskeget, Monomoy, Seal, and Green Islands) ranged from no change on Green Island to high rates of increase on the other three islands, with a maximum increase of 26.3% (95%CI: 21.6 - 31.4%; [5]). These high rates of increase provide further support for the hypothesis that seals from Canada are continually supplementing the breeding population in U.S. waters.

Strong evidence exists to suggest that interactions between right whales and both the fixed gear fisheries in the U.S. and Canada and vessel strikes in the U.S. are contributing substantially to the decline of the species [6]. Further, right whale distribution has changed since 2010. New research suggests that recent climate driven changes in ocean circulation have resulted in right whale distribution changes driven by increased warm water influx through the Northeast Channel, which has reduced the primary right whale prey (the copepod *Calanus finmarchicus*) in the central and eastern portions of the Gulf of Maine [6–8]. Additional potential stressors include offshore wind development, which overlaps with important habitat areas used year-round by right whales, including mother and calf migration corridors and foraging habitat [9,10]. This area is also the only known right whale winter foraging habitat. Additional information can be found in the [offshore wind risks section](#).

The UMEs are under investigation and are likely the result of multiple drivers. For the large whale UMEs, human interaction appears to have contributed to increased mortalities, although investigations are not complete. An investigation into the cause of the seal UME so far suggests avian flu virus as a potential cause.

A climate vulnerability assessment is currently underway for Atlantic and Gulf of Mexico marine mammal populations and will be reported on in future versions of this report.

Risks to meeting fishery management objectives

Climate and Ecosystem Productivity

Large scale climate related changes in the ecosystem can lead to changes in important habitats and ecological interactions, potentially resulting in regime shifts and ecosystem reorganization.

Climate Change Indicators: ocean temperature, heatwaves, currents, acidification

Ocean and estuarine temperature and salinity The ocean continues to warm, altering habitat conditions experienced by a wide range of species. 2022 was among the warmest years on record in the North Atlantic [11] and ocean temperatures continue to warm at both the surface (Fig. 28) and bottom (Fig. 29) throughout the Mid-Atlantic. Bottom temperature shows a long term warming trend in all seasons, while sea surface temperature shows significant long term warming in spring, summer, and fall. Seasonal sea surface temperatures in 2022 were above average for most of the year, however late spring storms caused deep mixing, which delayed stratification and surface warming in late spring and early summer.




Figure 28: MAB (grey outline) seasonal sea surface temperature (SST) time series overlaid onto 2021 seasonal spatial anomalies. Seasons are defined as: Jan-Mar for winter, Apr-Jun for spring, Jul-Sep for summer, and Oct-Dec for fall.




Figure 29: MAB seasonal bottom temperature (BT) anomaly time series. Seasons are defined as: Jan-Mar for winter, Apr-Jun for spring, Jul-Sep for summer, and Oct-Dec for fall. The final 2 years of each time series (open circles) are modeled estimates subject to change.

In addition to increasing temperatures overall, ocean summer conditions now last longer within each year. In the MAB, the transition date from warm stratified summer conditions to well mixed cool fall conditions is getting later (Fig. 30).




Figure 30: Ocean summer length in the MAB: the annual total number of days between the spring thermal transition date and the fall thermal transition date. The transition dates are defined as the day of the year when surface temperatures changeover from cool to warm conditions in the spring and back to cool conditions in the fall.

The Chesapeake Bay experienced a warmer-than-average winter 2022, and average conditions in the spring and summer. Fall 2022 was cooler relative to the baseline period 2008-2021 as measured by satellites¹³ and by buoys¹⁴ (Fig. 31, left panel), which also indicated above-average salinity in the Chesapeake Bay throughout the summer and fall (Fig. 31, right panel).




Figure 31: NOAA Chesapeake Bay Interpretive Buoy System Gooses Reef bouy sea water temperature (left) and salinity (right); Red = 2022, Blue = Long term average 2010-2020.

Extreme temperature events The increase in surface and bottom water temperature observed in the Northeast US may represent long term incremental stress on marine organisms, especially those relying on cooler water habitats for some or all life stages. In addition to changes in long-term average conditions, short-term extreme temperature events can produce acute stress on marine organisms, especially when the baseline temperature is increasing. To identify these extreme events separately from the baseline warming, we have changed our methods describing marine heatwaves (MHWs, [12]; [13]; [14]) to remove the global warming signal. Therefore, these indicators look different than in previous reports, but MHWs identified now are truly extreme departures from an already warming ecosystem. A combination of long-term ocean warming and MHWs should be used to assess total heat stress on marine organisms.

In 2022, the Mid-Atlantic Bight experienced two distinct surface marine heatwaves starting on August 29th and November 7th, lasting 9 and 11 days respectively (Fig. 32). Both ranked low among all recorded MWHs (75th and 73rd respectively). The top 4 strongest surface MHWs in the MAB occurred during the last ten years, with the two events in 2012 ranked as 1st and 3rd. No bottom MHWs were observed in 2022. The strongest bottom MHWs occurred in the fall of 1985 followed by the second strongest in the winter/spring of 2012.

¹³<https://coastwatch.noaa.gov/cw/index.html>

¹⁴<https://buoybay.noaa.gov/>




Figure 32: Marine heatwave events (red shading above black line) in the Mid-Atlantic occurring in 2022.

Ocean currents and features Variability of the Gulf Stream is one of the major drivers of changes in the oceanographic conditions of the Slope Sea and subsequently the Northeast U.S. continental shelf [15]. Changes in the Gulf Stream and Slope Sea can affect large-scale climate phenomena as well as local ecosystems and coastal communities. During the last decade, the Gulf Stream has become less stable and shifted northward [16,17] (Fig. 33). A more northern Gulf Stream position is associated with warmer ocean temperature on the northeast shelf [18], a higher proportion of Warm Slope Water in the Northeast Channel, and increased sea surface height along the U.S. east coast [19].




Figure 33: Index representing changes in the location of the Gulf Stream north wall. Positive values represent a more northerly Gulf Stream position.

Since 2008, the Gulf Stream has moved closer to the Grand Banks, reducing the supply of cold, fresh, and oxygen-rich Labrador Current waters to the Northwest Atlantic Shelf [20]. Nearly every year since 2010, warm slope water made up more than 75% of the annual slope water proportions entering the Gulf of Maine. In 2017 and 2019, almost no cooler Labrador Slope water entered the Gulf of Maine through the Northeast Channel (Fig. 34). The changing proportions of source water affect the temperature, salinity, and nutrient inputs to the Gulf of Maine ecosystem. In 2021, warm slope water continued to dominate (86.1%) inputs to the Gulf of Maine.




Figure 34: Proportion of Warm Slope Water (WSW) and Labrador Slope Water (LSW) entering the Gulf of Maine through the Northeast Channel.

The increased instability of the Gulf Stream position and warming of the Slope Sea may also be connected to the regime shift increase in the number of warm core rings formed annually in the Northwest Atlantic [15,21] (Fig. 35). When warm core rings and eddies interact with the continental slope they can transport warm, salty water to the continental shelf [22], which can alter the habitat and disrupt seasonal movements of fish [23]. Transport of offshore water onto the shelf is happening more frequently [23,24], and can contribute to marine heatwaves in the Mid-Atlantic Bight [22,25] as well as the movement of shelf-break species inshore [23,26,27].

2022 had the same number of warm core rings (21) as 2021, but most of the 2022 rings formed east of 60 W and fewer were observed near the shelf break region.




Figure 35: Warm core ring formation on the Northeast U.S. Shelf: Annual number of rings.

Changes in ocean temperature and circulation alter habitat features such as the seasonal cold pool, a 20–60 m thick band of cold, relatively uniform near-bottom water that persists from spring to fall over the mid and outer shelf of the MAB and southern flank of Georges Bank [28,29]. The cold pool plays an essential role in the structuring of the MAB ecosystem. It is a reservoir of nutrients that feeds phytoplankton productivity, is essential fish spawning and nursery habitat, and affects fish distribution and behavior [28,30]. The average temperature of the cold pool is getting warmer over time [31,32], the area is getting smaller [33], and the duration is getting shorter (Fig. 36).




Figure 36: Seasonal cold pool indices: mean temperature within the cold pool, cold pool persistence, and spatial extent.

Ocean Acidification Ocean acidification (OA) has caused measured declines in global ocean pH, and is projected to continue declining if high carbon dioxide emissions continue [34]. OA also changes the availability of minerals required by organisms to form calcified structures such as shells. Calcifying conditions in seawater can be determined by measuring aragonite saturation state (Ω_{Arag}), the tendency of a common type of calcium carbonate, aragonite, to form or dissolve. When Ω_{Arag} is less than 1, shells and other calcium carbonate structures begin to dissolve. Typical surface ocean Ω_{Arag} is 2-4, but extremes can be <1 or >5 [35]. As the ocean absorbs carbon dioxide, both pH and Ω_{Arag} decrease and can cause organisms to respond with reduced survival, calcification rates, growth, and reproduction, as well as impaired development, and/or changes in energy allocation [37]. However, sensitivity levels vary, and some organisms exhibit negative responses to calcification and other processes when Ω_{Arag} is as low as 3.

Summer-time (2007-present) Ω_{Arag} on the U.S. Northeast Shelf varies in space and time, ranging from 0.64 to 2.49 (Fig. 37, left panel). Spatially, the lowest bottom Ω_{Arag} has occurred in the Gulf of Maine, western Long Island Sound, nearshore to mid-shelf waters of the Mid-Atlantic Bight off the coast of New Jersey, and in waters > 1000 meters. Ω_{Arag} was at or below the sensitivity levels for both Atlantic sea scallop [38] and longfin squid [39,40] in Long Island Sound and the nearshore and mid-shelf regions of the New Jersey shelf (Fig. 37, right panels). The sensitivity levels of bottom Ω_{Arag} occurred during August 2016, July 2018, and August 2019 for both species, and additionally in August 2021 for the Atlantic sea scallop.




Figure 37: Left panel: Bottom aragonite saturation state (Ω_{Arag} ; summer only: June-August) on the U.S. Northeast Shelf based on quality-controlled vessel- and glider-based datasets from 2007-present. Right panel: Locations where summer bottom Ω_{Arag} were at or below the laboratory-derived sensitivity level for Atlantic sea scallop (top panel) and longfin squid (bottom). Gray circles indicate locations where carbonate chemistry samples were collected, but bottom Ω_{Arag} values were higher than sensitivity values determined for that species.

Ecosystem Productivity Indicators: phytoplankton, zooplankton, forage fish, fish condition

Phytoplankton Phytoplankton support the food web as the primary food source for zooplankton and filter feeders such as shellfish. Numerous environmental and oceanographic factors interact to drive the abundance, composition, spatial distribution, and productivity of phytoplankton. In 2022, MAB phytoplankton biomass (surface chlorophyll) was above average in winter, but below average in August and September. Below average phytoplankton biomass could be due to reduced nutrient flow to the surface and/or increased grazing pressure. Chlorophyll concentrations were above average in early fall and a fall bloom was detected in November/December. Primary productivity (the rate of photosynthesis) was average through spring, above average in the summer and average in the fall (Fig. 38).




Figure 38: Weekly chlorophyll concentrations and primary productivity in the Mid-Atlantic are shown by the colored line for 2022. The long-term mean is shown in black and shading indicates +/- 1 standard deviation.

The seasonal cycle of phytoplankton size distribution shows that the winter/spring and fall bloom periods are dominated by larger-celled microplankton, while smaller-celled nanoplankton dominate during the warmer summer months. The proportion of the smallest phytoplankton, picoplankton (0.2-2 microns), is relatively constant throughout the year. In 2022, microplankton proportions were average for most of the year, and above average peaks correspond to the bloom periods observed in chlorophyll concentration (Fig. 39).




Figure 39: The annual climatology (1998-2022) percent composition of the phytoplankton size classes in the Mid-Atlantic based on satellite observations in the shaded portions. The 2022 proportions for the microplankton (>20 microns, green) and nanoplankton (2-20 microns, orange) are shown in the bold lines.

Zooplankton The zooplankton community is changing in the MAB. Two dominant groups show long term trends: ‘sea butterflies’ (pteropods) show a long term increase in the MAB, and the copepod *Pseudocalanus* spp. has a long term decreasing trend (Fig. 40). Pteropods are important prey items for planktivores such as herring and mackerel,

as well as some sea birds. Despite being susceptible to shell degradation by ocean acidification, their abundance has remained above long term mean since 2004. Pseudocalanus spp. are important prey for many larval fish species, and can influence phytoplankton standing stock through grazing. Pseudocalanus spp. abundance has been below the long term mean since 2000 and continues to decrease with increasing temperature.




Figure 40: Abundance Anomalies of pseudocalanus and pteropods in Mid-Atlantic Bight.

Forage Fish Energy Content Nutritional value (energy content) of juvenile and adult forage fish as prey is related to environmental conditions, fish growth, and reproductive cycles. Forage energy density measurements from NEFSC trawl surveys 2017-2022 are building toward a time series to evaluate trends (Fig. 41). Data from the fall 2021 and spring 2022 survey measurements were consistent with previous reports: the energy density of Atlantic herring increased to over 7 kJ/g wet weight, but was still well below that observed in the 1980s and 1990s (10.6-9.4 kJ/ g wet weight). Silver hake, longfin squid (*Loligo* in figure) and shortfin squid (*Illex* in figure) remain lower than previous estimates [41,42]. Energy density of alewife, butterfish, sand lance, and Atlantic mackerel varies seasonally, with seasonal estimates both higher and lower than estimates from previous decades.




Figure 41: Forage fish energy density mean and standard deviation by season and year, compared with 1980s (solid line; Steimle and Terranova 1985) and 1990s (dashed line; Lawson et al. 1998) values.

Forage Fish Biomass Index The amount of forage fish available in the ecosystem combined with the energy content of the forage species determines the amount of energy potentially available to predators in the ecosystem. Changes in the forage base could pose a risk to managed and protected species production. A new spatially-explicit forage index estimated the combined biomass of 20 forage species using stomach contents information from 22 predatory fish species collected on bottom trawl surveys. While the resulting indices show no long term trends in the Mid-Atlantic, they do show overall higher forage fish in fall relative to spring (Fig. 42), with highest forage biomass during fall in the mid-1980s. Changes in the distribution of forage biomass also affects predator distribution. Spatial subsets of this index were included in the bluefish research track stock assessment to investigate forage-driven changes in bluefish availability to recreational fisheries and surveys.




Figure 42: Forage fish index based on spring and fall survey predator diets.

Fish Condition The health and well being of individual fish can be related to body shape condition indices (i.e., weight at a given length) such as relative condition index, which is the ratio of observed weight to predicted weight based on length [43]. Heavier and fatter fish at a given length have higher relative condition which is expected to improve growth, reproductive output, and survival. A pattern of generally good condition was observed across many MAB species prior to 2000, followed by a period of generally poor condition from 2001-2010, with a mix of good and poor condition from 2011-2019. Condition was again mixed in 2022, but a number of species improved in condition from the relatively low condition year in 2021 (Fig. 43). Preliminary results of synthetic analyses show that changes in temperature, zooplankton, fishing pressure, and population size influence the condition of different fish species.




Figure 43: Condition factor for fish species in the MAB based on fall NEFSC bottom trawl survey data. MAB data are missing for 2017 due to survey delays, and no survey was conducted in 2020.

Fish Productivity We describe patterns of aggregate fish productivity in the Mid-Atlantic with the small fish per large fish anomaly indicator, derived from NEFSC bottom trawl survey data (Fig. 44). The indicator shows that productivity has been declining in this region since 2010. A similar analysis based on stock assessment model outputs (recruitment per spawning stock biomass anomaly) for stocks primarily inhabiting the Mid-Atlantic region also shows a decline in productivity.




Figure 44: Fish productivity measures. Left: Small fish per large fish survey biomass anomaly in the Mid-Atlantic Bight. Right: assessment recruitment per spawning stock biomass anomaly for stocks mainly in the Mid-Atlantic. The summed anomaly across species is shown by the black line.

Ecosystem Structure Indicators: distribution shifts, diversity, predators

As noted in the Landings Implications section above, stocks are shifting distribution throughout the region. In aggregate, fish stocks are moving northeast along the shelf and into deeper waters.

Zooplankton diversity is increasing in the MAB, while adult fish diversity indices appear stable over time, with

current values within one standard deviation from most historic estimates (see [Diversity Indicators section, above](#)).

Indicators for shark populations, combined with information on gray seals (see [Protected Species Implications section, above](#)), suggests predator populations range from stable (sharks, Fig. 45) to increasing (seals) in the MAB. Stable predator populations suggest stable predation pressure on managed species, but increasing predator populations may reflect increasing predation pressure.




Figure 45: Estimated number of sharks per unit effort from Highly Migratory Species Pelagic Observer Program data.

Stock status is mixed for Atlantic Highly Migratory Species (HMS) stocks (including sharks, swordfish, billfish, and tunas) occurring in the Mid-Atlantic region. While there are several HMS species considered to be overfished or that have unknown stock status, the population status for some managed Atlantic sharks and tunas is at or above the biomass target (Fig. 46), suggesting the potential for robust predator populations among these managed species.




Figure 46: Summary of single species status for HMS stocks; key to species names at <https://noaa-edab.github.io/tech-doc/atlantic-highly-migratory-species-stock-status.html>.

As noted in the [Protected Species section](#), gray seal populations are increasing. Harbor and gray seals occupying New England waters are generalist predators that consume more than 30 different prey species. An evaluation of

hard parts found in seal stomachs showed that harbor and gray seals predominantly exploit abundant demersal fish species (i.e., red, white, and silver hake). Other relatively abundant prey species found in hard-part remains include sand lance, yellowtail flounder, four-spotted flounder, Gulf Stream flounder, haddock, herring, redfish, and squids.

A stable isotope study utilizing gray seal scat samples obtained from Massachusetts habitats showed individual gray seals can specialize on particular prey [44]. It also found that gray seals vary their diet seasonally, focusing on demersal inshore species prior to the spring molt, and offshore species such as sandlance after molting. DNA studies on gray seal diet in Gulf of Maine and Massachusetts waters found spiny dogfish and Jonah crab present in gray seal scat samples [45,46], with sandlance and menhaden dominant off Monomoy, MA [47]. Skate and crab remains were also found in gray seal stomach remains. In contrast to direct feeding, it is uncertain if the presence of skates and crabs is due to secondary consumption or scavenging.

Habitat Risk Indicators: habitat assessments, submerged aquatic vegetation, estuarine habitat quality, fishing gear impacts

Habitat Assessments The Northeast Regional Marine Fish Habitat Assessment (NRHA) is a collaborative effort to describe and characterize estuarine, coastal, and offshore fish habitat distribution, abundance, and quality in the Northeast. This includes mapping inshore and offshore habitat types used by focal fish species, summarizing impacts of habitat climate vulnerability on these species, modeling predicted future species distributions, and developing a publicly accessible decision support tool to visualize these results. This is a three-year project led by the New England and Mid-Atlantic Fishery Management Councils in collaboration with many partners including NOAA Fisheries¹⁵.

New habitat model-based richness estimates Species richness was derived from habitat models for 55 common species sampled by the spring and fall NEFSC bottom trawl surveys during the years 2000-2019 as part of the NRHA. The joint species distribution model controls for differences in capture efficiency across survey vessels, revealing patterns of declining richness in the Mid-Atlantic Bight and increasing richness in more northerly regions (i.e., the Gulf of Maine; Fig. 47). These patterns reflect the decreasing probability of occurrence of cooler-water species in the south (Atlantic cod, American plaice, pollock, thorny skate) and the growing prevalence of warm-water species in the north (weakfish, spotted hake, and black sea bass), likely as a result of rising water temperatures.




Figure 47: Habitat model-based species richness for 55 common species sampled by NEFSC bottom trawl surveys.

¹⁵<https://www.mafmc.org/nrha>

Submerged Aquatic Vegetation Submerged aquatic vegetation (SAV) is designated as a Habitat Area of Particular Concern (HAPC) for summer flounder and is important habitat for many fish species, particularly during vulnerable juvenile stages. Increased SAV coverage (including wild celery, water stargrass, and hydrilla) in the tidal fresh areas of the Chesapeake Bay (Fig. 48) has been attributed to restoration efforts. This ecosystem engineering has improved water quality, promoting further expansions of SAV meadows. However, in the higher salinity region near the mouth of the Chesapeake Bay (Fig. 48), increased water temperatures continue to inhibit eelgrass expansion. In 2021, the return to normal water temperature in the summer corresponded to a slight improvement in both eelgrass and widgeon grass coverage.




Figure 48: Submerged Aquatic Vegetation (SAV) coverage in tidal fresh and high salinity regions of the Chesapeake Bay.

Fishing Gear Impacts Estimates of the impacts of fishing gear on habitat are available through the habitat section of the Northeast Ocean Data Portal¹⁶. The data portal hosts selected outputs from the Northeast Fishing Effects Model which combines seafloor data (sediment type, energy regime) with fishing effort data to generate percent habitat disturbance estimates in space and time. More detailed information can be found in the Synthetic Indicator Catalog.¹⁷

Implications

Links between climate change and managed species Estuarine, nearshore, and offshore habitats support many life stages of state and federally managed species, and are highly vulnerable to climate change. Below we highlight how recently observed habitat changes affect several key managed species in Chesapeake Bay and in both nearshore and offshore waters of the MAB. Overall, multiple drivers interact differently for each species, producing a range of population impacts.

Estuarine habitat and managed species Relative habitat use of Chesapeake Bay by several finfish species, including Atlantic croaker, spot, summer flounder, weakfish, clearnose skate, and horseshoe crab is declining [48]. There is evidence suitable habitat for juvenile summer flounder growth has declined by 50% or more [49]. Climate change is expected to continue impacting habitat function and use for multiple species. Restoration of oyster reefs (see below) and marshes could help address these challenges.

Average water temperatures in 2022 (Fig. 31, left) and below-average hypoxic volume throughout the summer suggest favorable conditions for striped bass and blue crabs. Strong winds from the remnants of Hurricane Ian reduced hypoxia by mixing the water column in early October. However, the juvenile striped bass index was low, similar to the past four years, and the total population of blue crabs was at its lowest point in the history of the winter dredge survey. Lower winter temperatures may have contributed to higher overwintering mortality of adult female and juvenile blue crabs. The updated ASMFC striped bass stock assessment shows population numbers

¹⁶<https://www.northeastoceandata.org/data-explorer/>

¹⁷<https://noaa-edab.github.io/catalog/northeast-fishing-effects-model.html>

remain below the management threshold. Habitat conditions in the Chesapeake Bay could be one factor limiting striped bass population recovery and may have contributed to poor blue crab recruitment over the past few years, leading to lower overall abundances.

Forage and structure-forming species were likely favored by 2022 conditions in Chesapeake Bay. Average water temperatures in 2022 and above-average salinity conditions mean a suitable habitat year for bay anchovy, a key forage species. Bay anchovy abundances are directly correlated with the area of suitable habitat. Above-average salinities beginning in June 2022 (Fig.31, right) were associated with strong oyster recruitment [50]. However, oyster populations are severely depleted from historical levels. Large-scale restoration in 10 tributaries across the Chesapeake Bay is helping recover oyster reef habitat and populations in select areas.

Offshore habitat and managed species Ocean acidification also has different implications, depending on the species and life stage. Summer aragonite saturation was at or below the sensitivity levels for both Atlantic sea scallop and longfin squid in Long Island Sound and the nearshore and mid-shelf regions of the New Jersey shelf (Fig. 37, right panels) several times over the past decade. Recent lab studies have found that surf clams exhibited metabolic depression in a pH range of 7.46-7.28 [51]. Aggregated data from 2007-2021 show that summer bottom ocean pH (7.69-8.07) has not yet reached the metabolic depression threshold observed for surfclams in lab studies so far. The projected effects of changing temperature and ocean chemistry over the coming century may alter surfclam growth and reproduction [52].

While offshore habitat conditions have degraded for some species, they have improved for others. Between 2017 and 2021, extraordinarily high availability of northern shortfin squid (*Illex*) were observed in the Mid-Atlantic, resulting in high fishery catch per unit effort (CPUE) and early fishery closures. High instances of squid catch near the shelf break are significantly related to low bottom temperatures (< 10 degrees C), high salinity (>35.6 psu), increased chlorophyll frontal activity, as well as the presence and orientation of warm core rings. Warm core rings are an important contributor to squid availability, likely influencing habitat conditions across different life stages and as a transport mechanism of higher salinity water to the shelf. In addition, fishing effort is often concentrated on the eastern edge of warm core rings, which are associated with upwelling and enhanced productivity. There were fewer warm core rings near the continental shelf in 2022, which combined with economic fishery drivers may have contributed to total catch of *Illex* squid being 20% less than the total catch reported in 2021.

Marine heatwave impacts The adjustment to the marine heatwave methodology shows that extreme temperature events happen intermittently in many years, but have not been increasing over time in the Mid-Atlantic. While temperature variability in isolation has not changed, considering the overall increase in ocean temperature at both the surface and the bottom in the region, extreme events can represent additional stress to organisms. While marine heatwaves lasting over days may disturb the marine environment, long lasting events such as the warming in 2012 (Fig. 49) can have significant impacts to the ecosystem [25]. The 2012 heatwave affected the lobster fishery most notably, but other species also shifted their geographic distributions and seasonal cycles [53]. During the 2017 event, warm water fish typically found in the Gulf Stream were caught in shallow waters near Block Island, RI [23].

Mid-Atlantic Marine Heatwave Intensity




Figure 49: Marine heatwave maximum intensity (left) and total days each year (right) in the Mid-Atlantic Bight.

Cold pool impacts Changes in the cold pool habitat can affect species distribution, recruitment, and migration timing for multiple federally managed species. Southern New England-Mid Atlantic yellowtail flounder recruitment and settlement are related to the strength of the cold pool [31]. The settlement of pre-recruits during the cold pool event represents a bottleneck in yellowtail life history, during which a local and temporary increase in bottom temperature negatively impacts the survival of the settlers. Including the effect of cold pool variations on yellowtail recruitment reduced retrospective patterns and improved the skill of short-term forecasts in a stock assessment model [31,32]. The cold pool also provides habitat for the ocean quahog [33,54]. Growth rates of ocean quahogs in the MAB (southern portion of their range) have increased over the last 200 years whereas little to no change has been documented in the northern portion of their range in southern New England, likely a response to a warming and shrinking cold pool [55].

Distribution shift impacts Trends for a suite of 48 commercially or ecologically important fish species along the entire Northeast Shelf continue to show movement towards the northeast and generally into deeper water (Fig. 9). Habitat model-based species richness suggests shifts of both cooler and warmer water species to the northeast (Fig. 47). Similar patterns have been found for marine mammals, with multiple species shifting northeast between 2010 and 2017 in most seasons (Fig. 50, [56]).

Protected Species Distribution Shifts




Figure 50: Direction and magnitude of core habitat shifts, represented by the length of the line of the seasonal weighted centroid for species with more than 70 km difference between 2010 and 2017 (tip of arrow).

Shifting species distributions alter both species interactions and fishery interactions. In particular, shifting species distributions can alter expected management outcomes from spatial allocations and bycatch measures based on historical fish and protected species distributions.

Ecosystem productivity change impacts Climate and associated changes in the physical environment affect ecosystem productivity, with warming waters affecting the rate of photosynthesis at the base of the food web. Warm temperatures can increase the rate of primary production, however they also increase stratification, which limits the flux of deep water nutrients to the surface. Thus most of the increased summer production in the MAB is from smaller phytoplankton and may not translate into increased fish biomass.

While pteropods are increasing over time, smaller zooplankton are periodically shifting abundance between the larger, more nutritious copepod *Calanus finmarchicus* and smaller bodied copepods, and common *Pseudocalanus* copepods show a long term decrease in the MAB. The nutritional content of forage fish changes seasonally in response to ecosystem conditions, with apparent declines in energy density for Atlantic herring and *Illex* squid relative to the 1980s, but similar energy density for other forage species. Overall forage fish biomass has fluctuated in the MAB over time. Some of these factors are now being linked to the relative condition of managed fish.

The apparent decline in productivity across multiple managed species in the MAB, along with mixed fish conditions in 2022, also suggest changing ecosystem productivity at multiple levels. During the 1990s high relative abundance of smaller bodied copepods and a lower relative abundance of *Calanus finmarchicus* was associated with regime shifts to higher fish recruitment [57]. The unprecedented climate signals along with the trends toward lower productivity across multiple managed species indicate a need to continually evaluate whether management reference points remain appropriate, and to evaluate if ecosystem regime shifts have occurred or reorganization is in progress.

Other Ocean Uses: Offshore Wind

Indicators: development timeline, revenue in lease areas, coastal community vulnerability

As of January 2023, 31 offshore wind development projects are proposed for construction over the next decade in the Northeast (timelines and project data are based on Tables E-2, E-4, and E-4-2 of South Fork Wind Farm Final Environmental Impact Statement). Offshore wind areas are anticipated to cover 2.4 million acres by 2030 in the Greater Atlantic region (Fig. 51). Beyond 2030 values include acreage for future areas in the Central Atlantic and Gulf of Maine Area planning area for floating research array.





Figure 51: Proposed wind development on the northeast shelf.



Wind area boundaries, construction data and timelines are frequently updated. This map contains the most recent published information as of Dec 2022

Figure 52: All Northeast Project areas by year construction ends (each project has 2 year construction period).

Just over 3,400 foundations and more than 9,000 miles of inter-array and offshore export cables are proposed to

date. The colored chart in Fig. 52 also presents the offshore wind development timeline in the Greater Atlantic region with the estimated year that foundations would be constructed (matches the color of the wind areas). These timelines and data estimates are expected to shift but represent the most recent information available as of January 2023. Based on current timelines, the areas affected would be spread out such that it is unlikely that any one particular area would experience full development at one time. Future wind development areas are also presented. Additional call areas, which may eventually become lease areas, totalling over 488,000 acres in the Central Atlantic¹⁸ may be identified for BOEM’s anticipated 2023 lease sale. It’s anticipated that the Central Atlantic leases will fulfill outstanding offshore wind energy production goals for VA and NC.

Based on federal vessel logbook data, commercial fishery revenue from trips in the current offshore wind lease areas and the draft Central Atlantic Bight Primary and Secondary Call Areas have varied annually from 2008-2021, with less than \$1 million in revenue overlapping with these areas for most fisheries. However, some fisheries see periodic spikes in revenue overlap with wind energy lease areas, including up to \$4.7 million affected in the surfclam fishery and nearly \$4.3 million affected in the longfin squid fishery in 2008 and 2016, respectively.(Fig. 53).




Figure 53: Fishery revenue in wind energy lease areas in the Mid-Atlantic.

Of MAFMC-managed fisheries, the chub mackerel fishery could be the fishery most affected by offshore wind development, with a maximum of 17% of annual regional fishery revenue occurring within potential wind lease areas and the Central Atlantic draft call areas during this period, followed by the surfclam (16%), black sea bass (15%), ocean quahog (13%), and blueline tilefish fisheries (10%). The spiny dogfish fishery was the least affected, at 3% maximum annual revenue affected, while 5% of annual revenues were affected for several others (bluefish, butterfish, and summer flounder). A maximum of 10% of the annual longfin squid revenues were affected by these areas, with similar effects for the scup (9%), Atlantic mackerel (8%), monkfish (7%) and golden tilefish (6%) fisheries (see Table 2). While up to 14% of annual *Illex* squid revenue overlapped with offshore wind areas, this is likely overestimated due to the precision of logbook data when compared to vessel monitoring system data (see Table 2).

Table 2: Top Species Landings and Revenue from Wind Energy Areas. * Landings and revenue for these species are likely underestimated due to limited coverage of these fisheries in historic reporting requirements for vessels issued federal permits by the NMFS Greater Atlantic Regional Fisheries Office. However, such limitations also suggest an inaccurately higher proportion of such landings and revenues in existing lease areas. ** Clearnose skates were reported separately from skates, which is presumed to include all skates managed under the Northeast skate complex. *** Based on comparison with other data sources, the high values for *Illex* squid are likely overestimates affected by the methods used to model logbook data to estimate spatial overlap of fishign operations with wind energy areas.

NEFMC, MAFMC, and ASMFC Managed Species	Maximum Percent Total Annual Regional Species Landings	Maximum Percent Total Annual Regional Species Revenue
Black drum*	36	34
American eel*	15	29
Clearnose skate**	19	20

¹⁸https://www.boem.gov/sites/default/files/images/draft_wea_primary_secondary3.jpg

NEFMC, MAFMC, and ASMFC Managed Species	Maximum Percent Total Annual Regional Species Landings	Maximum Percent Total Annual Regional Species Revenue
Atlantic menhaden*	25	19
Atlantic chub mackerel*	16	17
Atlantic surfclam	17	16
Black sea bass	15	15
Yellowtail flounder	15	15
Illex squid***	14	14
Offshore hake	14	14
Ocean quahog	13	13
Atlantic sea scallops	13	12
Blueline tilefish*	8	10
Skates**	10	10
Longfin squid	9	9
Scup	8	9
Atlantic mackerel	8	8
Monkfish	9	7
Red hake	11	7

Proposed wind development areas interact with the region’s federal scientific surveys. Scientific surveys are impacted by offshore wind in four ways: 1. Exclusion of NOAA Fisheries’ sampling platforms from the wind development area due to operational and safety limitations; 2.Impacts on the random-stratified statistical design that is the basis for scientific assessments, advice, and analyses; 3.Alteration of benthic and pelagic habitats, and airspace in and around the wind energy development, requiring new designs and methods to sample new habitats; and, 4.Reduced sampling productivity through navigation impacts of wind energy infrastructure on aerial and vessel survey operations. Increase vessel transit between stations may decrease data collections that are already limited by annual days-at-sea day allocations. The total survey area overlap ranges from 1-14% for all Greater Atlantic federal surveys. Individual survey strata have significant interaction with wind, including the sea scallop survey (up to 96% of individual strata) and the bottom trawl survey (BTS, up to 60% strata overlap). Additionally, up to 50% of the southern New England North Atlantic right whale survey’s area overlaps with proposed project areas. A region-wide survey mitigation program is underway [58].

Equity and environmental justice (EJ) are priority concerns with offshore wind development and fisheries impacts in the Northeast. Fig. 54 links historic port revenue (2008-2021) from within all wind lease areas as a proportion of the port’s total revenue based on vessel trip reports as described in the revenue and landings of species in the wind indicator above. The range (minimum and maximum) of total percent revenue from within wind energy areas is presented in the graph and Mid-Atlantic ports are sorted from greatest to least revenue from within wind areas.

For example, Atlantic City, NJ had a minimum of 11% and maximum of 30% overlap of fisheries revenue in potential wind development areas to the total port fisheries revenue between 2008-2021. Those communities that score Med-High or higher in at least one of the vulnerability indicators that address environmental justice concerns (i.e., Poverty, Population Composition, Personal Disruption; see indicator definitions) are noted with a triangle. Gentrification pressure is also highlighted here, with those communities that score Med-High or higher in one or more gentrification pressure indicators (i.e., Housing Disruption, Retiree Migration, Urban Sprawl) represented with a circle (Fig. 54). BOEM reports that cumulative offshore wind development (if all proposed projects are developed) could have moderate impacts on low-income members of environmental justice communities who work in the commercial fishing and for-hire fishing industry due to disruptions to fish populations, restrictions on navigation and increased vessel traffic, as well as existing vulnerabilities of low-income workers to economic impacts [59].




Figure 54: Percent of Mid-Atlantic port revenue from Wind Energy Areas (WEA) in descending order from most to least port revenue from WEA. EJ = Environmental Justice.

Some ports in New England land Mid-Atlantic managed species from wind areas as well. For the maximum percent value reported in each New England port, the majority (at least 50% based on both value and pounds) of those landings were Mid-Atlantic managed species within wind areas for Barnstable, MA, Boston, MA, Hyannis, MA, North Kingstown/Davisville, RI, and Point Judith, RI. Woods Hole, MA would be added to this list based on pounds only, but did not exceed 50% of value from Mid-Atlantic managed species within wind areas.




Figure 55: Percent of New England port revenue with majority MAFMC landings from Wind Energy Areas (WEA) in descending order from most to least port revenue from WEA. EJ = Environmental Justice.

Top fishing communities high in environmental justice concerns (i.e., Atlantic City, NJ, Newport News, VA, Hobucken and Beaufort, NC) should be considered in decision making to reduce the social and economic impacts and aid in the resilience and adaptive capacity of underserved communities. It also highlights communities where we need to provide further resources to reach underserved and underrepresented groups and create opportunities for and directly involve these groups in the decision-making process.

Implications

Current plans for rapid buildout of offshore wind in a patchwork of areas spreads the impacts differentially throughout the region (Fig. 52).

Up to 17% of maximum annual fisheries revenue for major Mid-Atlantic commercial species in lease areas and draft call areas could be forgone or reduced and associated effort displaced if all sites are developed. Displaced fishing effort can alter historic fishing area, timing, and method patterns, which can in turn change habitat, species (managed and protected), and fleet interactions. Several factors, including fishery regulations, fishery availability, and user conflicts affect where, when, and how fishing effort may be displaced, along with impacts to and responses of affected fish species.

Planned development overlaps right whale mother and calf migration corridors and a significant foraging habitat that is used throughout the year [9] (Fig 56). Turbine presence and extraction of energy from the system could alter local oceanography [60] and may affect right whale prey availability. For example, persistent foraging hotspots of right whales and seabirds overlap on Nantucket Shoals, where unique hydrography aggregates enhanced prey densities [61,62]. Wind leases (OCS-A 0521 and OCS-A 0522) currently intersect these hotspots on the southwestern corner of Nantucket Shoals and a prominent tidal front associated with invertebrate prey swarms important to seabirds and possibly right whales. Proposed wind development areas also bring increased vessel strike risk to whales from construction and operation vessels, in addition to potential impacts such as displacement, increased levels of communication masking, and elevated stress hormones from pile driving and operational noise.




Figure 56: Northern Right Whale persistent hotspots and Wind Energy Areas.

Scientific data collection surveys for ocean and ecosystem conditions, fish, and protected species will be altered, potentially increasing uncertainty for stock assessments and associated management decision making.

The increase of offshore wind development can have both positive (e.g., employment opportunities) and negative (e.g., space-use conflicts) effects. Continued increase in coastal development and gentrification pressure has resulted in loss of fishing infrastructure space within ports. Understanding these existing pressures can allow for avoiding and mitigating negative impacts to our shore support industry and communities dependent on fishing. Some of the communities with the highest fisheries revenue overlap with offshore wind development areas that are also vulnerable to gentrification pressure are Point Pleasant and Atlantic City, NJ, Ocean City, MD, and Beaufort, NC.

Contributors

Editors (NOAA NMFS Northeast Fisheries Science Center, NEFSC): Sarah Gaichas, Kimberly Bastille, Geret DePiper, Kimberly Hyde, Scott Large, Sean Lucey, Laurel Smith

Contributors (NEFSC unless otherwise noted): Kimberly Bastille, Aaron Beaver (Anchor QEA), Andy Beet, Ruth Boettcher (Virginia Department of Game and Inland Fisheries), Mandy Bromilow and CJ Pellerin (NOAA Chesapeake Bay Office), Joseph Caracappa, Baoshan Chen (Stony Brook University), Doug Christel (GARFO), Patricia Clay, Lisa Colburn, Jennifer Cudney and Tobey Curtis (NMFS Atlantic HMS Management Division), Geret DePiper, Dan Dorfman (NOAA-NOS-NCCOS), Hubert du Pontavice, Emily Farr and Grace Roskar (NMFS Office of Habitat Conservation), Michael Fogarty, Paula Fratantoni, Kevin Friedland, Marjy Friedrichs (VIMS),

Sarah Gaichas, Ben Galuardi (GAFRO), Avijit Gangopadhyay (School for Marine Science and Technology, University of Massachusetts Dartmouth), James Gartland (Virginia Institute of Marine Science), Lori Garzio (Rutgers University), Glen Gawarkiewicz (Woods Hole Oceanographic Institution), Sean Hardison, Kimberly Hyde, John Kosik, Steve Kress and Don Lyons (National Audubon Society's Seabird Restoration Program), Young-Oh Kwon and Zhuomin Chen (Woods Hole Oceanographic Institution), Andrew Lipsky, Sean Lucey, Chris Melrose, Shannon Meseck, Ryan Morse, Ray Mroch (SEFSC), Brandon Muffley (MAFMC), Kimberly Murray, Janet Nye (University of North Carolina at Chapel Hill), Chris Orphanides, Richard Pace, Debi Palka, Tom Parham (Maryland DNR), Charles Perretti, Grace Saba and Emily Slesinger (Rutgers University), Vincent Saba, Sarah Salois, Chris Schillaci (GARFO), Amy Schueller (SEFSC), Teresa Schwemmer (Stony Brook University), Dave Secor (CBL), Angela Silva, Adrienne Silver (UMass/SMASST), Laurel Smith, Talya ten Brink (GARFO), Bruce Vogt (NOAA Chesapeake Bay Office), Ron Vogel (University of Maryland Cooperative Institute for Satellite Earth System Studies and NOAA/NESDIS Center for Satellite Applications and Research), John Walden, Harvey Walsh, Changhua Weng, Timothy White (Environmental Studies Program, BOEM), Mark Wuenschel

Document Orientation

The figure format is illustrated in Fig 57a. Trend lines are shown when slope is significantly different from 0 at the $p < 0.05$ level. An orange line signifies an overall positive trend, and purple signifies a negative trend. To minimize bias introduced by small sample size, no trend is fit for < 30 year time series. Dashed lines represent mean values of time series unless the indicator is an anomaly, in which case the dashed line is equal to 0. Shaded regions indicate the past ten years. If there are no new data for 2022, the shaded region will still cover this time period. The spatial scale of indicators is either coastwide, Mid-Atlantic states (New York, New Jersey, Delaware, Maryland, Virginia, North Carolina), or at the Mid-Atlantic Bight (MAB) Ecosystem Production Unit (EPU, Fig. 57b) level.




Figure 57: Document orientation. a. Key to figures. b. The Northeast Large Marine Ecosystem.

Fish and invertebrates are aggregated into similar feeding categories (Table 3) to evaluate ecosystem level trends in predators and prey.

Table 3: Feeding guilds and management bodies.

Guild	MAFMC	Joint	NEFMC	State or Other
Apex Predator				bluefin tuna, shark uncl, swordfish, yellowfin tuna
Piscivore	bluefish, longfin squid, northern shortfin squid, summer flounder	goosefish, spiny dogfish	acadian redfish, atlantic cod, atlantic halibut, clearnose skate, little skate, offshore hake, pollock, red hake, silver hake, smooth skate, thorny skate, white hake, winter skate	fourspot flounder, john dory, sea raven, striped bass, weakfish, windowpane
Planktivore	atlantic mackerel, butterfish		atlantic herring	alewife, american shad, blackbelly rosefish, blueback herring, cusk, longhorn sculpin, lumpfish, menhaden, northern sand lance, northern searobin, sculpin uncl
Benthivore	black sea bass, scup, tilefish		american plaice, barndoor skate, crab, red deepsea, haddock, ocean pout, rosette skate, winter flounder, witch flounder, yellowtail flounder	american lobster, atlantic wolffish, blue crab, cancer crab uncl, chain dogfish, cunner, jonah crab, lady crab, smooth dogfish, spider crab uncl, squid cuttlefish and octopod uncl, striped searobin, tautog
Benthos	atlantic surfclam, ocean quahog		sea scallop	blue mussel, channeled whelk, sea cucumber, sea urchin and sand dollar uncl, sea urchins, snails(conchs)

References

1. Gaichas SK, DePiper GS, Seagraves RJ, Muffley BW, Sabo M, Colburn LL, et al. Implementing Ecosystem Approaches to Fishery Management: Risk Assessment in the US Mid-Atlantic. *Frontiers in Marine Science*. 2018;5. doi:[10.3389/fmars.2018.00442](https://doi.org/10.3389/fmars.2018.00442)
2. Friedland KD, Langan JA, Large SI, Selden RL, Link JS, Watson RA, et al. Changes in higher trophic level productivity, diversity and niche space in a rapidly warming continental shelf ecosystem. *Science of The Total Environment*. 2020;704: 135270. doi:[10.1016/j.scitotenv.2019.135270](https://doi.org/10.1016/j.scitotenv.2019.135270)
3. Thunberg EM. Northeast Region Fisheries Impacts from COVID-19. US Seafood Industry and For-Hire Sector Impacts from COVID-19: 2020 in Perspective NOAA Tech Memo NMFS-SPO-221. 2021. pp. 53–64. Available: <https://spo.nmfs.noaa.gov/sites/default/files/TM221.pdf>
4. Pace RM, Williams R, Kraus SD, Knowlton AR, Pettis HM. Cryptic mortality of North Atlantic right whales. *Conservation Science and Practice*. 2021;n/a: e346. doi:<https://doi.org/10.1111/csp.2346>
5. Wood SA, Murray KT, Josephson E, Gilbert J. Rates of increase in gray seal (*Halichoerus grypus atlantica*) pupping at recolonized sites in the United States, 1988–2019. Swanson B, editor. *Journal of Mammalogy*. 2020;101: 121–128. doi:[10.1093/jmammal/gyz184](https://doi.org/10.1093/jmammal/gyz184)
6. Hayes S, Gardner S, Garrison LP, Henry A, Leandro L. North Atlantic Right Whales-Evaluating Their Recovery Challenges in 2018. NOAA Tech Memo NMFS NEFSC 247. 2018.
7. Record N, Runge J, Pendleton D, Balch W, Davies K, Pershing A, et al. Rapid Climate-Driven Circulation Changes Threaten Conservation of Endangered North Atlantic Right Whales. *Oceanography*. 2019;32. doi:[10.5670/oceanog.2019.201](https://doi.org/10.5670/oceanog.2019.201)
8. Sorochan KA, Plourde S, Morse R, Pepin P, Runge J, Thompson C, et al. North Atlantic right whale (*Eubalaena glacialis*) and its food: (II) interannual variations in biomass of *Calanus* spp. On western North Atlantic shelves. *Journal of Plankton Research*. 2019;41: 687–708. doi:[10.1093/plankt/fbz044](https://doi.org/10.1093/plankt/fbz044)
9. Quintana-Rizzo E, Leiter S, Cole TVN, Hagbloom MN, Knowlton AR, Nagelkirk P, et al. Residency, demographics, and movement patterns of North Atlantic right whales *Eubalaena glacialis* in an offshore wind energy development area in southern New England, USA. *Endangered Species Research*. 2021;45: 251–268. doi:[10.3354/esr01137](https://doi.org/10.3354/esr01137)
10. Schick RS, Halpin PN, Read AJ, Slay CK, Kraus SD, Mate BR, et al. Striking the right balance in right whale conservation. *Canadian Journal of Fisheries and Aquatic Sciences*. 2009;66: 1399–1403. doi:[10.1139/F09-115](https://doi.org/10.1139/F09-115)
11. Cheng L, Abraham J, Trenberth KE, Fasullo J, Boyer T, Mann ME, et al. Another Year of Record Heat for the Oceans. *Advances in Atmospheric Sciences*. 2023; doi:[10.1007/s00376-023-2385-2](https://doi.org/10.1007/s00376-023-2385-2)
12. Hobday AJ, Alexander LV, Perkins SE, Smale DA, Straub SC, Oliver ECJ, et al. A hierarchical approach to defining marine heatwaves. *Progress in Oceanography*. 2016;141: 227–238. doi:[10.1016/j.pocean.2015.12.014](https://doi.org/10.1016/j.pocean.2015.12.014)
13. Jacox MG, Alexander MA, Bograd SJ, Scott JD. Thermal displacement by marine heatwaves. *Nature*. 2020;584: 82–86. doi:[10.1038/s41586-020-2534-z](https://doi.org/10.1038/s41586-020-2534-z)
14. Jacox MG, Alexander MA, Amaya D, Becker E, Bograd SJ, Brodie S, et al. Global seasonal forecasts of marine heatwaves. *Nature*. 2022;604: 486–490. doi:[10.1038/s41586-022-04573-9](https://doi.org/10.1038/s41586-022-04573-9)
15. Gangopadhyay A, Gawarkiewicz G, Silva ENS, Silver AM, Monim M, Clark J. A Census of the Warm-Core Rings of the Gulf Stream: 1980–2017. *Journal of Geophysical Research: Oceans*. 2020;125: e2019JC016033. doi:[10.1029/2019JC016033](https://doi.org/10.1029/2019JC016033)
16. Andres M. On the recent destabilization of the Gulf Stream path downstream of Cape Hatteras. *Geophysical Research Letters*. 2016;43: 9836–9842. doi:[10.1002/2016GL069966](https://doi.org/10.1002/2016GL069966)
17. Caesar L, Rahmstorf S, Robinson A, Feulner G, Saba V. Observed fingerprint of a weakening Atlantic Ocean overturning circulation. *Nature*. 2018;556: 191–196. doi:[10.1038/s41586-018-0006-5](https://doi.org/10.1038/s41586-018-0006-5)
18. Zhang R, Vallis GK. The Role of Bottom Vortex Stretching on the Path of the North Atlantic Western Boundary Current and on the Northern Recirculation Gyre. *Journal of Physical Oceanography*. 2007;37: 2053–2080. doi:[10.1175/JPO3102.1](https://doi.org/10.1175/JPO3102.1)
19. Goddard PB, Yin J, Griffies SM, Zhang S. An extreme event of sea-level rise along the Northeast coast of North America in 2009–2010. *Nature Communications*. 2015;6. doi:[10.1038/ncomms7346](https://doi.org/10.1038/ncomms7346)

20. Gonçalves Neto A, Langan JA, Palter JB. Changes in the Gulf Stream preceded rapid warming of the Northwest Atlantic Shelf. *Communications Earth & Environment*. 2021;2: 1–10. doi:[10.1038/s43247-021-00143-5](https://doi.org/10.1038/s43247-021-00143-5)
21. Gangopadhyay A, Gawarkiewicz G, Silva ENS, Monim M, Clark J. An Observed Regime Shift in the Formation of Warm Core Rings from the Gulf Stream. *Scientific Reports*. 2019;9: 1–9. doi:[10.1038/s41598-019-48661-9](https://doi.org/10.1038/s41598-019-48661-9)
22. Chen K, Gawarkiewicz G, Yang J. Mesoscale and Submesoscale Shelf-Ocean Exchanges Initialize an Advective Marine Heatwave. *Journal of Geophysical Research: Oceans*. 2022;127: e2021JC017927. doi:<https://doi.org/10.1029/2021JC017927>
23. Gawarkiewicz G, Todd R, Zhang W, Partida J, Gangopadhyay A, Monim M-U-H, et al. The Changing Nature of Shelf-Break Exchange Revealed by the OOI Pioneer Array. *Oceanography*. 2018;31: 60–70. doi:[10.5670/oceanog.2018.110](https://doi.org/10.5670/oceanog.2018.110)
24. Gawarkiewicz G, Fratantoni P, Bahr F, Ellertson A. Increasing Frequency of Mid-depth Salinity Maximum Intrusions in the Middle Atlantic Bight. *Journal of Geophysical Research: Oceans*.
25. Gawarkiewicz G, Chen K, Forsyth J, Bahr F, Mercer AM, Ellertson A, et al. Characteristics of an Advective Marine Heatwave in the Middle Atlantic Bight in Early 2017. *Frontiers in Marine Science*. 2019;6. Available: <https://www.frontiersin.org/article/10.3389/fmars.2019.00712>
26. Potter IF, Galuardi B, Howell WH. Horizontal movement of ocean sunfish, *Mola mola*, in the northwest Atlantic. *Marine Biology*. 2011;158: 531–540. doi:[10.1007/s00227-010-1578-2](https://doi.org/10.1007/s00227-010-1578-2)
27. Worm B, Lotze HK, Myers RA. Predator diversity hotspots in the blue ocean. *Proceedings of the National Academy of Sciences*. 2003;100: 9884–9888. doi:[10.1073/pnas.1333941100](https://doi.org/10.1073/pnas.1333941100)
28. Lentz SJ. Seasonal warming of the Middle Atlantic Bight Cold Pool. *Journal of Geophysical Research: Oceans*. 2017;122: 941–954. doi:[10.1002/2016JC012201](https://doi.org/10.1002/2016JC012201)
29. Chen Z, Curchitser E, Chant R, Kang D. Seasonal Variability of the Cold Pool Over the Mid-Atlantic Bight Continental Shelf. *Journal of Geophysical Research: Oceans*. 2018;123: 8203–8226. doi:[10.1029/2018JC014148](https://doi.org/10.1029/2018JC014148)
30. Miles T, Murphy S, Kohut J, Borsetti S, Munroe D. Offshore Wind Energy and the Mid-Atlantic Cold Pool: A Review of Potential Interactions. *Marine Technology Society Journal*. 2021;55: 72–87. doi:[10.4031/MTSJ.55.4.8](https://doi.org/10.4031/MTSJ.55.4.8)
31. Miller TJ, Hare JA, Alade LA. A state-space approach to incorporating environmental effects on recruitment in an age-structured assessment model with an application to southern New England yellowtail flounder. *Canadian Journal of Fisheries and Aquatic Sciences*. 2016;73: 1261–1270. doi:[10.1139/cjfas-2015-0339](https://doi.org/10.1139/cjfas-2015-0339)
32. Pontavice H du, Miller TJ, Stock BC, Chen Z, Saba VS. Incorporating environmental effects from ocean models improves a marine fish stock assessment. *ICES Journal of Marine Science*.
33. Friedland KD, Miles T, Goode AG, Powell EN, Brady DC. The Middle Atlantic Bight Cold Pool is warming and shrinking: Indices from in situ autumn seafloor temperatures. *Fisheries Oceanography*. 2022;31: 217–223. doi:[10.1111/fog.12573](https://doi.org/10.1111/fog.12573)
34. Intergovernmental Panel on Climate Change (IPCC), editor. Technical Summary. *The Ocean and Cryosphere in a Changing Climate: Special Report of the Intergovernmental Panel on Climate Change*. Cambridge: Cambridge University Press; 2022. pp. 39–70. doi:[10.1017/9781009157964.002](https://doi.org/10.1017/9781009157964.002)
35. Jiang L-Q, Feely RA, Carter BR, Greeley DJ, Gledhill DK, Arzayus KM. Climatological distribution of aragonite saturation state in the global oceans. *Global Biogeochemical Cycles*. 2015;29: 1656–1673. doi:[10.1002/2015GB005198](https://doi.org/10.1002/2015GB005198)
36. Kroeker KJ, Kordas RL, Crim R, Hendriks IE, Ramajo L, Singh GS, et al. Impacts of ocean acidification on marine organisms: Quantifying sensitivities and interaction with warming. *Global Change Biology*. 2013;19: 1884–1896. doi:[10.1111/gcb.12179](https://doi.org/10.1111/gcb.12179)
37. Saba GK, Goldsmith KA, Cooley SR, Grosse D, Meseck SL, Miller AW, et al. Recommended priorities for research on ecological impacts of ocean and coastal acidification in the U.S. Mid-Atlantic. *Estuarine, Coastal and Shelf Science*. 2019;225: 106188. doi:[10.1016/j.ecss.2019.04.022](https://doi.org/10.1016/j.ecss.2019.04.022)

38. Cameron LP, Grabowski JH, Ries JB. Effects of elevated pCO₂ and temperature on the calcification rate, survival, extrapallial fluid chemistry, and respiration of the Atlantic Sea scallop *Placopecten magellanicus*. *Limnology and Oceanography*. 2022;67: 1670–1686. doi:[10.1002/lno.12153](https://doi.org/10.1002/lno.12153)
39. Zakroff C, Mooney TA, Berumen ML. Dose-dependence and small-scale variability in responses to ocean acidification during squid, *Doryteuthis pealeii*, development. *Marine Biology*. 2019;166: 62. doi:[10.1007/s00227-019-3510-8](https://doi.org/10.1007/s00227-019-3510-8)
40. Zakroff CJ, Mooney TA. Antagonistic Interactions and Clutch-Dependent Sensitivity Induce Variable Responses to Ocean Acidification and Warming in Squid (*Doryteuthis pealeii*) Embryos and Paralarvae. *Frontiers in Physiology*. 2020;11. Available: <https://www.frontiersin.org/articles/10.3389/fphys.2020.00501>
41. Steimle F, Terranova R. Energy Equivalents of Marine Organisms from the Continental Shelf of the Temperate Northwest Atlantic. *Journal of Northwest Atlantic Fishery Science*. 1985;6. doi:[10.2960/J.v6.a11](https://doi.org/10.2960/J.v6.a11)
42. Lawson JW, Magalhães AM, Miller EH. Important prey species of marine vertebrate predators in the north-west Atlantic: Proximate composition and energy density. *Marine Ecology Progress Series*. 1998;164: 13–20. Available: <https://www.jstor.org/stable/24825521>
43. Le Cren ED. The Length-Weight Relationship and Seasonal Cycle in Gonad Weight and Condition in the Perch (*Perca fluviatilis*). *Journal of Animal Ecology*. 1951;20: 201–219. doi:[10.2307/1540](https://doi.org/10.2307/1540)
44. Hernandez KM, Bogomolni AL, Moxley JH, Waring GT, DiGiovanni RA, Hammill MO, et al. Seasonal variability and individual consistency in gray seal (*Halichoerus grypus*) isotopic niches. *Canadian Journal of Zoology*. 2019;97: 1071–1077. doi:[10.1139/cjz-2019-0032](https://doi.org/10.1139/cjz-2019-0032)
45. Ono KA, Steinbeiser CM, Coco AB, Sheehan MJ, Beck AJ, Dufault MN, et al. Detecting spiny dogfish in grey seal diets. *Conservation Genetics Resources*. 2019;11: 481–485. doi:[10.1007/s12686-018-1044-x](https://doi.org/10.1007/s12686-018-1044-x)
46. McCosker C, Flanders K, Ono K, Dufault M, Mellone D, Olson Z. Metabarcoding Fecal DNA Reveals Extent of *Halichoerus grypus* (Gray Seal) Foraging on Invertebrates and Incidence of Parasite Exposure. *Northeastern Naturalist*. 2020;27: 681–700. doi:[10.1656/045.027.0409](https://doi.org/10.1656/045.027.0409)
47. Flanders KR, Olson ZH, Ono KA. Utilizing next-generation sequencing to identify prey DNA in western North Atlantic grey seal *Halichoerus grypus* diet. *Marine Ecology Progress Series*. 2020;655: 227–240. doi:[10.3354/meps13520](https://doi.org/10.3354/meps13520)
48. Schonfeld AJ, Gartland J, Latour RJ. Spatial differences in estuarine utilization by seasonally resident species in Mid-Atlantic Bight, USA. *Fisheries Oceanography*. 2022;31: 615–628. doi:[10.1111/fog.12611](https://doi.org/10.1111/fog.12611)
49. Fabrizio M, Tuckey T, Smith S, Ross P, Snyder R, Wang H, et al. Characterization of Nursery Habitats used by Black Sea Bass and Summer Flounder in Chesapeake Bay and the Coastal Lagoons. *Reports*. 2022; doi:[10.25773/PJCC-RG41](https://doi.org/10.25773/PJCC-RG41)
50. Kimmel DG, Tarnowski M, Newell RIE. The Relationship between Interannual Climate Variability and Juvenile Eastern Oyster Abundance at a Regional Scale in Chesapeake Bay. *North American Journal of Fisheries Management*. 2014;34: 1–15. doi:[10.1080/02755947.2013.830999](https://doi.org/10.1080/02755947.2013.830999)
51. Pousse E, Poach ME, Redman DH, Sennefelder G, White LE, Lindsay JM, et al. Energetic response of Atlantic surfclam *Spisula solidissima* to ocean acidification. *Marine Pollution Bulletin*. 2020;161: 111740. doi:[10.1016/j.marpolbul.2020.111740](https://doi.org/10.1016/j.marpolbul.2020.111740)
52. Pousse É, Munroe D, Hart D, Hennen D, Cameron LP, Rheuban JE, et al. Dynamic energy budget modeling of Atlantic surfclam, *Spisula solidissima*, under future ocean acidification and warming. *Marine Environmental Research*. 2022;177: 105602. doi:[10.1016/j.marenvres.2022.105602](https://doi.org/10.1016/j.marenvres.2022.105602)
53. Mills K, Pershing A, Brown C, Chen Y, Chiang F-S, Holland D, et al. Fisheries Management in a Changing Climate: Lessons From the 2012 Ocean Heat Wave in the Northwest Atlantic. *Oceanography*. 2013;26. doi:[10.5670/oceanog.2013.27](https://doi.org/10.5670/oceanog.2013.27)
54. Powell EN, Ewing AM, Kuykendall KM. Ocean quahogs (*Arctica islandica*) and Atlantic surfclams (*Spisula solidissima*) on the Mid-Atlantic Bight continental shelf and Georges Bank: The death assemblage as a recorder of climate change and the reorganization of the continental shelf benthos. *Palaeogeography, Palaeoclimatology, Palaeoecology*. 2020;537: 109205. doi:[10.1016/j.palaeo.2019.05.027](https://doi.org/10.1016/j.palaeo.2019.05.027)

55. Pace SM, Powell EN, Mann R. Two-hundred year record of increasing growth rates for ocean quahogs (*Arctica islandica*) from the northwestern Atlantic Ocean. *Journal of Experimental Marine Biology and Ecology*. 2018;503: 8–22. doi:[10.1016/j.jembe.2018.01.010](https://doi.org/10.1016/j.jembe.2018.01.010)
56. Chavez-Rosales S, Josephson E, Palka D, Garrison L. Detection of Habitat Shifts of Cetacean Species: A Comparison Between 2010 and 2017 Habitat Suitability Conditions in the Northwest Atlantic Ocean. *Frontiers in Marine Science*. 2022;9. Available: <https://www.frontiersin.org/articles/10.3389/fmars.2022.877580>
57. Perretti C, Fogarty M, Friedland K, Hare J, Lucey S, McBride R, et al. Regime shifts in fish recruitment on the Northeast US Continental Shelf. *Marine Ecology Progress Series*. 2017;574: 1–11. doi:[10.3354/meps12183](https://doi.org/10.3354/meps12183)
58. Hare JA, Blythe BJ, Ford KH, Godfrey-McKee S, Hooker BR, Jensen BM, et al. NOAA Fisheries and BOEM Federal Survey Mitigation Implementation Strategy - Northeast U.S. Region. *Northeast Fisheries Science Center (U.S.)*, editor. 2022; doi:[10.25923/jqse-x746](https://doi.org/10.25923/jqse-x746)
59. BOEM. Vineyard Wind 1 Offshore Wind Energy Project Supplement to the Draft Environmental Impact Statement. OCS EIS/EA, BOEM 2020-025 [Internet]. 2020. Available: <https://www.boem.gov/sites/default/files/documents/renewable-energy/Vineyard-Wind-1-Supplement-to-EIS.pdf>
60. Christiansen N, Daewel U, Djath B, Schrum C. Emergence of Large-Scale Hydrodynamic Structures Due to Atmospheric Offshore Wind Farm Wakes. *Frontiers in Marine Science*. 2022;9. Available: <https://www.frontiersin.org/article/10.3389/fmars.2022.818501>
61. White TP, Veit RR. Spatial ecology of long-tailed ducks and white-winged scoters wintering on nantucket shoals. *Ecosphere*. 2020;11: e03002. doi:<https://doi.org/10.1002/ecs2.3002>
62. Sorochan KA, Plourde S, Baumgartner MF, Johnson CL. Availability, supply, and aggregation of prey (*Calanus* spp.) in foraging areas of the North Atlantic right whale (*Eubalaena glacialis*). *ICES Journal of Marine Science*. 2021;78: 3498–3520. doi:[10.1093/icesjms/fsab200](https://doi.org/10.1093/icesjms/fsab200)